arXiv:1209.3444v1 [math.AG] 15 Sep 2012

Comparison theorems for deformation functors

Jan Arthur Christophersen Jan O. Kleppe

September 18, 2012

Abstract

This is a preliminary report on our study of comparisons of deformations of algebras to deformations of schemes via invariant theory. We generalize comparison theorems of the second author for projective schemes. We consider deformation functors for a scheme X which is a good quotient of a quasi-affine scheme X' by a linearly reductive group G and compare them to invariant deformations of any affine G-scheme containing X' as an open invariant subset.

Introduction

Given a projective scheme X defined by equations $f_1, \dots, f_m \in k[x_0, \dots, x_n]$, perturbing the equations in a flat manner so that they remain homogeneous induce deformations of X. In practice this is often the only way to construct examples of deformations. In more stringent terms we have a map between the degree 0 embedded deformations of the affine cone $C(X)$ and deformations of X in $\mathbb{P}^n$. If we take into account trivial deformations we get a map to the deformations of X as scheme. The question is when do we get all deformations this way.

To see precisely what is going on we should compare deformation functors on Artin rings. If $R = k[x_0, \dots, x_n]$ and $S = R/(f_1, \dots, f_m)$ then the above describes maps $\mathrm{Def}^0_{S/R} \to \mathrm{Hilb}_{X/\mathbb{P}^n}$ where $\mathrm{Def}^0_{S/R}$ is the functor of degree 0 deformations of S as R-algebra and $\mathrm{Def}^0_S \to \mathrm{Def}_X$ where Def^0_S is the functor of degree 0 deformations of S as k-algebra. In [Kle79] the second author gave exact conditions for when these maps are isomorphisms. The object of this paper is to generalize these to other situations where one compare can deformations of algebras to deformations of schemes.

The comparison map for projective schemes factors through deformations of the open subset of $C(X)$ where the vertex $\{0\}$ is removed. Thereafter one compares deformations to $X = (C(X) \setminus \{0\})/k^*$ via the quotient map. A natural question is if this can be generalized to closed subschemes of toric varieties corresponding to ideals in the Cox ring. It turns out one can go even further, i.e. the quotient need not be by a quasi-torus.

In this paper we consider schemes X which are good quotients of a quasi-affine scheme X' by a linearly reductive group G. We assume that $X' \subseteq \operatorname{Spec} S$ and that G acts on

S inducing the action on X'. We can then compare Def_S^G to Def_X where Def_S^G is the functor of invariant deformations of S. If this situation is embedded in another one we can compare with the local Hilbert functor as well. The precise definitions of these settings are formulated with what we call G-ferry quadruples - see Definition 2.1 and Definition 3.1. This generality allows us to say something about affine schemes like quotient singularities as well.

Linearly reductive groups have many properties coming from the Reynolds operator which make it possible to prove things, e.g. taking invariants is exact. Another reason to work with them is that the functor of invariant deformations is well defined and has the usual nice properties of a good deformation theory. This was proven by Rim in [Rim80].

Our main result on the local Hilbert functor is Theorem 3.4. The conditions are depth conditions along the complement of X' in $\operatorname{Spec} S$ and along the locus where the quotient map is not a G-bundle. We state also corollaries for subschemes of toric varieties and weighted projective space.

For the abstract deformation functor Def_X the results are not as exact due to the presence of infinitesimal automorphisms. It is not clear what the correct assumptions should be but we found it useful to use results of Altmann regarding rigidity of $\mathbb{Q}$-Gorenstein toric singularities as a guide. We get depth conditions as above but also along the locus where the isotropy groups are not finite. An important ingredient are what we call a set of Euler derivations coming from the Lie algebra of G. This was explained to us by Dmitry Timashev and made it possible to work with general groups and not just tori which we had originally studied.

The main results for Def_X are Theorem 4.10 and in the Cohen-Macaulay case Theorem 4.13. We conclude with applications to rigidity questions for toric varieties. All though we here only reprove known results the techniques seem to promise new applications.

As stated in the abstract this is a preliminary version. In a future version we believe we can weaken the assumptions somewhat and describe more applications.

Terminology. The functor of deformations of a scheme X or R-algebra S are denoted Def_X and $\mathrm{Def}_{S/R}$, a superscript 0 means of degree 0 in the graded case and a G means invariant. The functor of embedded deformations of a scheme $X \subset Y$ is denoted $\mathrm{Hilb}_{X/Y}$ and called the local Hilbert functor.

We use throughout the standard criteria for when a map of deformation functors is smooth, namely that it is surjective on tangent spaces and injective on obstruction spaces. When the tangent map is an isomorphism we say the map of deformation functors is an isomorphism. If the functors have hulls then the hulls in this case will be isomorphic.

Acknowledgments. We are grateful to Dmitry Timashev, Arne B Sletsøe and Nathan Owen Ilten for helpful discussions and answering questions. We would in particular thank Dmitry Timashev for explaining aspects of invariant theory that led to the the correct setting for our results.

1 Preliminaries

1.1 Cotangent cohomology statements

To fix notation we give a short description of the cotangent modules and sheaves. Given a ring R and an R-algebra S there is a complex of free S modules; the *cotangent complex* $\mathbb{L}_{\bullet}^{S/R}$. See e.g. [And74, p. 34] for a definition. For an S module M we get the *cotangent cohomology* modules $T^i(S/R; M) = H^i(\operatorname{Hom}_S(\mathbb{L}_{\cdot}^{S/R}, M))$. If R is the ground field we abbreviate $T^i(S/R; M) = T^i_S(M)$ and $T^i_S(S) = T^i_S = T^i_X$ if $X = \operatorname{Spec} S$.

If X is a scheme we may globalise these as follows. If $\mathcal{R}$ is a sheaf of rings and $\mathcal{S}$ an $\mathcal{R}$ algebra we set $\mathcal{L}_{\cdot}^{\mathcal{S}/\mathcal{R}}$ to be the complex of sheaves associated with the presheaves $U \mapsto \mathbb{L}_{\cdot}^{\mathcal{S}(U)/\mathcal{R}(U)}$. Let $\mathcal{F}$ be an $\mathcal{S}$ module. We get the cotangent cohomology sheaves $\mathcal{T}^i(\mathcal{S}/\mathcal{R}; \mathcal{F})$ as the cohomology sheaves of $\mathcal{H}om_{\mathcal{S}}(\mathcal{L}_{\cdot}^{\mathcal{S}/\mathcal{R}}, \mathcal{F})$ and the cotangent cohomology groups $T^i(\mathcal{S}/\mathcal{R}; \mathcal{F})$ as the cohomology of $\operatorname{Hom}_{\mathcal{S}}(\mathcal{L}_{\cdot}^{\mathcal{S}/\mathcal{R}}, \mathcal{F})$.

Because of the functoriality of these constructions we have $\mathcal{T}^i(\mathcal{S}/\mathcal{R}; \mathcal{F})$ as the sheaf associated to the presheaf $U \mapsto T^i(\mathcal{S}(U)/\mathcal{R}(U); \mathcal{F}(U))$ and $T^{\bullet}(\mathcal{S}/\mathcal{R}; \mathcal{F})$ as the hypercohomology of $\mathcal{H}om_{\mathcal{S}}(\mathcal{L}_{\cdot}^{\mathcal{S}/\mathcal{R}}, \mathcal{F})$. In particular there is a "local-global" spectral sequence

$$H^p(X, \mathcal{T}^i(\mathcal{S}/\mathcal{R}; \mathcal{F})) \Rightarrow T^n(\mathcal{S}/\mathcal{R}; \mathcal{F}) \,. \tag{1.1}$$

If $\mathcal{S}$ is the structure sheaf $\mathcal{O}_X$ and R is the ground field, then we abbreviate as above to $T^i_X(\mathcal{F})$.

The properties of the cotangent cohomology we use, e.g. the Zariski-Jacobi sequence and flat base change results, may be found in [And74]. We include here one result that does not seem to be well known. Let $Z \subseteq X$ be a closed subscheme, then following Laudal [Lau79, 3.2.10] one may define cotangent cohomology with support in Z denoted $T^i_Z(\mathcal{O}_X/\mathcal{R}; \mathcal{F})$. If $Z \subseteq X = \operatorname{Spec} S$ is given by $V(I)$, we write $T^i_I(S/R; S) = T^i_Z(\mathcal{O}_X/\mathcal{O}_R; \mathcal{O}_X)$.

Theorem 1.1. [Lau79, Theorem 3.2.11] *There is a long exact sequence*

$$\cdots \to T^i_Z(\mathcal{O}_X/\mathcal{R}; \mathcal{F}) \to T^i(\mathcal{O}_X/\mathcal{R}; \mathcal{F}) \to T^i(\mathcal{O}_{X\setminus Z}/\mathcal{R}; \mathcal{F}) \to T^{i+1}_Z(\mathcal{O}_X/\mathcal{R}; \mathcal{F}) \to \dots$$

and a spectral sequence yielding

$$T^p(\mathcal{O}_X/\mathcal{R}; \mathcal{H}^q_Z(\mathcal{F})) \Rightarrow T^n_Z(\mathcal{O}_X/\mathcal{R}; \mathcal{F}) \,.$$

1.2 Lemmata

We collect here some general statements in invariant theory and commutative algebra which we could not find a reference for.

Lemma 1.2. *Let G be a linearly reductive group acting on a noetherian k-algebra A, M an AG-module of finite type and N an AG-module. Let $F_{\bullet}$ be an AG projective resolution of M and $K^{\bullet}$ the induced subcomplex*

$$K^{\bullet} = \{\varphi \in \operatorname{Hom}_A(F_{\bullet}, N)^G : \varphi(F_{\bullet}^G) = 0\} \subseteq \operatorname{Hom}_A(F_{\bullet}, N) \,.$$

Then there is a long exact sequence of A^G-modules

$$\cdots \to H^i(K^\bullet) \to \operatorname{Ext}^i_A(M,N)^G \to \operatorname{Ext}^i_{A^G}(M^G,N^G) \to H^{i+1}(K^\bullet) \to \cdots .$$

Proof. Since taking G invariants is an exact functor, it is enough to prove that there is an exact sequence of complexes

$$0 \to K^\bullet \to \operatorname{Hom}_A(F_\bullet,N)^G \to \operatorname{Hom}_{A^G}(F_\bullet^G,N^G) \to 0$$

where the right map is $\varphi \mapsto \varphi_{|F_\bullet^G}$, which makes sense since φ is invariant. Clearly $K^\bullet$ is the kernel of this map. To prove that the right map is surjective consider the diagram

$$\begin{tikzcd}
F_\bullet \arrow[r, dashed, "\tilde\varphi"] \arrow[d, "\rho"] & N \arrow[d, two heads, "\rho"] \\
F_\bullet^G \arrow[r, "\varphi"] & N^G
\end{tikzcd}$$

where ρ is the Reynolds operator projecting onto the invariant subset. Since the $F_\bullet$ are projective the map $\varphi \circ \rho$ lifts to $\tilde\varphi$. ☐

Lemma 1.3. *Let G be a linearly reductive group acting on an n-dimensional k vector space V. Let $A = \operatorname{Sym}(V)$ and let N be an AG-module with $N^G = 0$. Then $\operatorname{Ext}^1_A(\Omega^1_{A/A^G},N)^G = 0$.*

Proof. Consider the cotangent sequence

$$\Omega^1_{A^G} \otimes_{A^G} A \to \Omega^1_A \to \Omega^1_{A/A^G} \to 0\,.$$

Now as AG module $\Omega^1_A \simeq A \otimes_k V^*$ where V^* is the dual vector space. If

$$A^G = k[\varphi_1,\dots,\varphi_N] \subseteq A = k[x_1,\dots,x_n]$$

then the beginning of an AG projective resolution of Ω^1_{A/A^G} looks like

$$\cdots \to A^N \xrightarrow{\left(\frac{\partial\varphi_j}{\partial x_i}\right)} A \otimes_k V^* \to \Omega^1_{A/A^G} \to 0\,.$$

Since of course A^N is generated by invariants Lemma 1.2 tells us that $\operatorname{Ext}^1_A(\Omega^1_{A/A^G},N)^G \to \operatorname{Ext}^1_{A^G}((\Omega^1_{A/A^G})^G,N^G)$ is injective. ☐

Lemma 1.4. *If A is a Cohen-Macaulay ring and $I \subset A$ a radical ideal of codimension d let*

$$J = \bigcap_{\substack{\mathfrak{p}\in \operatorname{Ass} I \\ \operatorname{codim}\mathfrak{p}=d}} \mathfrak{p}\,.$$

Then $H^d_J(A) \simeq H^d_I(A)$.

Proof. Assume $\mathfrak{p}_0 \in \operatorname{Ass} I = \{\mathfrak{p}_0, \ldots, \mathfrak{p}_n\}$, a set of distinct prime ideals, and $\operatorname{codim} \mathfrak{p}_0 > d$. Let $I' = \cap_{i \geq 1} \mathfrak{p}_i$. The interesting part of the Mayer-Vietoris sequence is

$$\cdots \to H^d_{\mathfrak{p}_0+I'}(A) \to H^d_{\mathfrak{p}_0}(A) \oplus H^d_{I'}(A) \to H^d_I(A) \to H^{d+1}_{\mathfrak{p}_0+I'}(A) \to \cdots .$$

Since A is Cohen-Macaulay $\operatorname{depth}_{\mathfrak{p}_0} A = \operatorname{codim} \mathfrak{p}_0 \geq d+1$. If $\operatorname{codim} \mathfrak{p}_0 + I' = d+1$ then we must also have $\operatorname{codim} \mathfrak{p}_0 = d+1$. Since $\mathfrak{p}_0$ is prime this implies that $I' \subseteq \mathfrak{p}_0$, but I is reduced so $\operatorname{Ass} I$ consists of minimal primes and we have contradiction. Thus we get an isomorphism $H^d_{I'}(A) \simeq H^d_I(A)$. In this manner we may remove all prime ideals of codimension strictly greater than d. ☐

Lemma 1.5. *If A is a Cohen-Macaulay integral domain, $I \subset A$ an ideal of codimension d contained in an ideal J of codimension strictly greater than d, then $H^0_J(H^d_I(A)) = 0$.*

Proof. We must exhibit an $x \in J$ which is not a zero divisor on $H^d_I(A)$. We may assume both I and J are reduced and by Lemma 1.4 that all associated primes of I are of codimension d. We clam there is an $x \in J$ such that $(I : x) = I$. Now

$$(I : x) = \bigcap_{\mathfrak{p} \in \operatorname{Ass} I} (\mathfrak{p} : x)$$

so we must find an x not in any of the associated primes of I. If this was not possible, then by the Prime Avoidance Lemma, J is contained in an associated prime of I. But then $\operatorname{codim} J \leq d$ which contradicts the assumption.

We have an exact sequence

$$0 \to H^{d-1}_I(A/(x)) \to H^d_I(A) \xrightarrow{\cdot x} H^d_I(A)\,.$$

Since $A/(x)$ is also Cohen-Macaulay we are done if we can prove that $\operatorname{codim} I/xI = d$ in $A/(x)$. This is true because x is a non-zero-divisor for A/I so $\dim A - \dim A/I = \dim A/x - \dim A/(I + (x))$. ☐

2 G-ferry quadruples

Definition 2.1. Let G be a linearly reductive algebraic group. Consider a quadruple (X, Z, S, J) where X is a noetherian k-scheme, Z is a (possibly empty) closed subset, S is a finitely generated k-algebra on which G acts and $J \subseteq S$ is a G invariant radical ideal. Set $X' = \operatorname{Spec} S \setminus V(J)$ and $U = X \setminus Z$. We say that (X, Z, S, J) is a *G-ferry quadruple* if there is a commutative diagram

$$\begin{tikzcd}
U' \arrow[r, hook] \arrow[d, "\pi|_{U'}"] & X' \arrow[r, hook] \arrow[d, "\pi"] & \operatorname{Spec} S \\
U \arrow[r, hook] & X &
\end{tikzcd}$$

where $U' = \pi^{-1}(U)$ and

- π is a good quotient of X' by the action G
- $\pi_{|U'}$ is a principal G bundle, in particular smooth
- $\operatorname{depth}_J S \geq 1$.

The sheaf of algebras $\mathcal{S} = \pi_* \mathcal{O}_{X'}$ is called *the ferry.*

We will throughout the rest of this paper use the above notation to refer to the various objects in the definition. Here are some examples.

Example 2.2. *The* Spec *construction.* If $X = \operatorname{Spec} A$ and G is trivial then $(X, \emptyset, A, (1))$ is a ferry quadruple.

Example 2.3. *The usual* Proj *construction.* If $X = \operatorname{Proj} A$ for A finitely generated $\mathbb{Z}$-graded k-algebra generated in degree 1 with irrelevant ideal $\mathfrak{m}$ and $G = k^*$ then $(X, \emptyset, A, \mathfrak{m})$ is a G-ferry quadruple.

Example 2.4. *The* Proj *construction in general.* Let S be a finitely generated $\mathbb{Z}_+$ graded k-algebra with $S_0 = k$ and set $X = \operatorname{Proj} S$. Assume $S = R/I$ where $R = k[x_0, \ldots, x_n]$ is graded with $\deg x_i = q_i \in \mathbb{N}$ and for $\mathfrak{m} = (x_0, \ldots, x_n)$, $\operatorname{depth}_{\mathfrak{m}} S \geq 1$. This defines an embedding of X into the weighted projective space $\mathbb{P}(\mathbf{q}) = \mathbb{P}(q_0, \ldots, q_n)$.

Following Miles Reid we say that $\mathbb{P}(\mathbf{q})$ is *well formed* if no $n-1$ of the $q_0, \ldots, q_n$ have a common factor. Similarly we will say that the corresponding grading on S is well formed. We may assume that all gradings are well formed, see e.g. [Del75, Proposition 1.3] or [Dol82, 1.3.1].) Let J_k be the ideal of R generated by $\{x_i : k \nmid q_i\}$ and set

$$J = \bigcap_{k \geq 2} J_k = \bigcap_{\substack{p \text{ prime} \\ p | \operatorname{lcm}(q_0, \ldots, q_n)}} J_p \,.$$

The singular locus of the well formed $\mathbb{P}(\mathbf{q})$ is $Z = V(J) \subseteq \mathbb{P}(\mathbf{q})$.

Then $(X, X \cap Z, S, \mathfrak{m})$ is a G-ferry quadruple.

Example 2.5. *The Cox construction for complex toric varieties.* This is our main example and it includes the previous ones, see e.g. [Cox95] and [CLS11, Chapter 5]. To fix notation for the rest of this paper we recall the construction. Using the standard notation for toric geometry let $X = X_\Sigma$ be an n-dimensional toric variety given by a fan Σ in $N_{\mathbb{R}}$. We assume here for simplicity that X_Σ has no torus factors, but this is not necessary for applying our results (see [CLS11, 5.1.11]). Let $\Sigma(1) = \{\rho_1, \ldots, \rho_N\}$ be the set of rays of Σ and let v_i be the primitive generator of $\rho_i \cap N$. The divisor class group of X is given by the exact sequence

$$0 \to \mathbb{Z}^n \xrightarrow{b} \mathbb{Z}^N \to \operatorname{Cl}(X) \to 0$$

where $b(u) = (< u, v_1 >, \ldots, < u, v_N >)$. The ring $S = \mathbb{C}[x_\rho : \rho \in \Sigma(1)]$ is naturally graded by the abelian group $\operatorname{Cl}(X)$ and with this grading it is called the *Cox ring* or *total homogeneous coordinate ring* of X.

For each cone σ in Σ there is a monomial

$$\prod_{\rho_i \not\subseteq \sigma} x_i \in S$$

and define $B(\Sigma)$ to be the ideal of S generated by these. It is called the *irrelevant ideal* of the Cox ring. Let $Z(\Sigma) = V(B(\Sigma)) \subseteq \operatorname{Spec} S$. Let G be the quasi-torus $\operatorname{Hom}_{\mathbb{Z}}(\operatorname{Cl}(X_\Sigma), \mathbb{C}^*)$. The theorem of Cox states that X is an almost geometric quotient for the action of G on $\operatorname{Spec} S \setminus Z(\Sigma)$.

Let $\operatorname{Sing}(X_\Sigma)$ be the singular locus, i.e.

$$\operatorname{Sing}(X_\Sigma) = \bigcup_{\substack{\sigma\in\Sigma \\ \sigma \text{ not smooth}}} V(\sigma)\,.$$

The smooth locus is given by the subfan consisting of smooth cones in Σ. Then it follows from the construction (see e.g. [CLS11, Exercise 5.1.10]) that $(X_\Sigma, \operatorname{Sing}(X_\Sigma), S, B(\Sigma))$ is a G-ferry quadruple. Note that a G invariant S-module M is the same thing as a $\operatorname{Cl}(X)$ graded S-module and that $M^G = M_0$, the degree $0 \in \operatorname{Cl}(X)$ part of M. Moreover if $\alpha = [D] \in \operatorname{Cl}(X)$ then $S_\alpha = \mathcal{O}_X(D)$. In particular the ferry in this case is

$$\mathcal{S} = \bigoplus_{[D]\in\operatorname{Cl}(X)} \mathcal{O}_X(D)\,.$$

Example 2.6. *Multigraded sheaves of algebras.* All the examples above and also the Cox ring construction for general varieties fit into the following set up. Let Γ be a finitely generated abelian group. Consider a reduced noetherian k-scheme X with a sheaf of Γ-graded algebras $\mathcal{S}$ which is locally of finite type. Assume $\mathcal{S}_0 = \mathcal{O}_X$ and that $S = H^0(X, \mathcal{S})$ is a finitely generated k-algebra. Then $X' = \operatorname{Spec}_X \mathcal{S} \subseteq \operatorname{Spec} S$ is quasi-affine. Set J to be the radical ideal defining $\operatorname{Spec} S \setminus X'$.

Let $G = \operatorname{Spec} k[\Gamma]$ be the corresponding quasi-torus and assume k and G are so that G is linearly reductive. Choose $Z \subset X$ so that G acts freely on $\pi^{-1}(X \setminus Z)$. (In the Cox ring construction one could choose $Z = \operatorname{Sing}(X)$.) Then (X, Z, S, J) is a G-ferry quadruple.

Example 2.7. *Quotient singularities.* Let $G \subset \operatorname{GL}_n(\mathbb{C})$ be a finite group without pseudo-reflections. Then $(\mathbb{C}^n/G, \operatorname{Sing}(\mathbb{C}^n/G), \mathcal{O}_{\mathbb{C}^n}, (1))$ is a G-ferry quadruple.

Example 2.8. *Homogeneous spaces.* Let G be a linearly reductive algebraic group and H a closed subgroup. Let S be the coordinate ring of the affine scheme G. Then $(G/H, \emptyset, S, (1))$ is a G-ferry quadruple.

3 Deformations of the embedded scheme - $\operatorname{Hilb}_{X/Y}$

3.1 G-subferry quadruples

Definition 3.1. Given a G-ferry quadruple (Y, W, R, J) and a G-invariant ideal $I \subseteq R$ contained in J we may construct a new G-ferry quadruple as follows. Let $S = R/I$, set

$$\begin{tikzcd}
 & & \operatorname{Spec} S \arrow[rr, hook] & & \operatorname{Spec} R \\
 & X' \arrow[ur, hook] \arrow[rr, hook, "f'"] \arrow[dd] & & Y' \arrow[ur, hook] \arrow[dd, "\pi"] & \\
U'_X \arrow[ur, hook] \arrow[rr, hook] \arrow[dd] & & U'_Y \arrow[ur, hook] \arrow[dd, "\pi_{|U'}"] & & \\
 & X \arrow[rr, hook, "f"] & & Y & \\
U_X \arrow[ur, hook] \arrow[rr, hook] & & U_Y \arrow[ur, hook] & &
\end{tikzcd}$$

Figure 1: The diagram for G-subferries.

$\bar{J} = J/I$ and assume $\operatorname{depth}_{\bar{J}} S \geq 1$. Let $\pi : Y' \to Y$ be the good quotient as in Definition 2.1. The quasi-affine scheme $X' = \operatorname{Spec} S \setminus V(J)$ is a closed G-invariant subset of Y'. Thus $\pi : X' \to X = \pi(X')$ is a good quotient where X has structure sheaf $(\pi_* \mathcal{O}_{X'})^G$. It follows that $(X, X \cap W, S, J/I)$ is also a G-ferry quadruple.We call it the *G-subferry quadruple induced by I.*

A G-subferry quadruple determines a diagram as in Figure 1 where $Z = X \cap W$, $U_X = X \cap U_Y$ and by definition $U'_X = \pi^{-1}(U_X)$. Let $\mathcal{S} = \pi_* \mathcal{O}_{X'}$ be the ferry of $(X, Z, S, \bar{J})$. Let $f : X \to Y$ be the closed embedding. Let I_W be the radical ideal of $\operatorname{Spec} R \setminus U'_Y$ and set $I_Z = (I_W + I)/I$. We will use this notation and the notation in the diagram throughout.

Now $\pi_{|U'_Y}$ is a principal bundle, so in particular inverse images of points are G-orbits. Thus for any G-invariant subset $V \subseteq U'_Y$ we have $\pi^{-1}(\pi(V)) = V$. Thus $X' \cap U'_Y = \pi^{-1}(\pi(X' \cap U'_Y)) = \pi^{-1}(\pi(X') \cap \pi(U'_Y))$ again since $x, y \in U'_Y$ and $\pi(x) = \pi(y)$ implies x and y are in the same G-orbit. This shows that $U'_X = X' \cap U'_Y$, so it is indeed an open subset of X' and the above diagram is commutative. Note that the front vertical square is Cartesian but the back one need not be.

3.2 Results

We begin with a general lemma.

Lemma 3.2. *If in a Cartesian square of schemes*

$$\begin{tikzcd}
X' \arrow[r, "f'"] \arrow[d] & Y' \arrow[d, "\pi"] \\
X \arrow[r, "f"] & Y
\end{tikzcd}$$

the morphism π is flat and affine and f, f' are closed immersions then

$$T^i(\mathcal{O}_X/f^{-1}\mathcal{O}_Y;\pi_*\mathcal{O}_{X'}) \simeq T^i(\mathcal{O}_{X'}/f'^{-1}\mathcal{O}_{Y'};\mathcal{O}_{X'})$$

for all $i \geq 0$.

Proof. We first compare the corresponding sheafs. Since all morphisms are affine and π is flat the diagram locally corresponds to a cocartesian diagram of rings

$$\begin{tikzcd}
A' & B' \arrow[l] \\
A \arrow[u] & B \arrow[l] \arrow[u, "\pi^\#"']
\end{tikzcd}$$

with $\pi^\#$ flat, so $T^i(A/B;A') \simeq T^i(A'/B';A')$. By comparing values on open affine sets we get therefore

$$\mathcal{T}^q(\mathcal{O}_X/f^{-1}\mathcal{O}_Y;\pi_*\mathcal{O}_{X'}) \simeq \pi_*\mathcal{T}^q(\mathcal{O}_{X'}/f'^{-1}\mathcal{O}_{Y'};\mathcal{O}_{X'})$$

for all $q \geq 0$. Since π is affine and being affine is preserved under base change, $H^p(X,\pi_*\mathcal{F}) \simeq H^p(X',\mathcal{F})$ for any quasi-coherent $\mathcal{F}$. Thus all terms in the two spectral sequences (1.1) for the two cohomology groups are isomorphic and the result follows. ☐

Lemma 3.3. *If* $\operatorname{depth}_Z \mathcal{S} \geq 2$ *then*

(i) $\operatorname{depth}_Z \mathcal{O}_X \geq 2$

(ii) $H^0_{\bar{J}}(S) = H^0_{I_Z}(S) = 0$

(iii) *there is an isomorphism* $H^1_{\bar{J}}(S) \simeq H^1_{I_Z}(S)$.

Proof. Since taking invariants is exact the first statement follows from $\mathcal{H}^i_Z(\mathcal{S})^G = \mathcal{H}^i_Z(\mathcal{O}_X)$. We have $\operatorname{depth}_{\bar{J}} S \geq 1$ by the assumption that $(X,Z,S,\bar{J})$ is a G-subferry, so we get (ii) and (iii) by considering the commutative diagram

$$\begin{tikzcd}
 & 0 \arrow[r] & S \arrow[r] \arrow[d, equal] & H^0(X,\mathcal{S}) \arrow[r] \arrow[d] & H^1_{\bar{J}}(S) \arrow[r] & 0 \\
0 \arrow[r] & H^0_{I_Z}(S) \arrow[r] & S \arrow[r] & H^0(U_X,\mathcal{S}_{|U_X}) \arrow[r] & H^1_{I_Z}(S) \arrow[r] & 0
\end{tikzcd}$$

where the middle vertical map is an isomorphism by the depth condition. ☐

Theorem 3.4. *If $(X,Z,S,\bar{J})$ is a G-subferry quadruple of (Y,W,R,J) induced by $I \subseteq R$ with ferry $\mathcal{S}$ and*

(i) $\operatorname{depth}_Z \mathcal{S} \geq 2$

(ii) $\operatorname{Hom}_R(I, H^1_{\bar{J}}(S))^G = 0$

then $\mathrm{Def}^G_{S/R}$ *and* $\mathrm{Hilb}_{X/Y}$ *are isomorphic deformation functors.*

Proof. We will prove that both deformation functors are isomorphic to Hilb_{U_X/U_Y}. Note first that if $f : X \to Y$ is any closed embedding then $T^0(\mathcal{O}_X/f^{-1}\mathcal{O}_Y;\mathcal{F}) = 0$ for any $\mathcal{O}_X$-module $\mathcal{F}$. Thus the spectral sequence in Theorem 1.1 yields $T^1_{I_Z}(S/R;S) \simeq \mathrm{Hom}_R(I, H^0_{I_Z}(S)) = 0$ by Lemma 3.3. Furthermore this and Lemma 3.3 (iii) shows that $T^2_{I_Z}(S/R;S) \simeq \mathrm{Hom}_R(I, H^1_{I_Z}(S)) \simeq \mathrm{Hom}_R(I, H^1_{\bar{J}}(S))$.

We start with applying the long exact sequence in Theorem 1.1 and get

$$0 \to T^1(S/R;S) \to T^1(\mathcal{O}_{U'_X}/R;\mathcal{O}_{U'_X}) \to T^2_{I_Z}(S/R;S) \to T^2(S/R;S) \to T^2(\mathcal{O}_{U'_X}/R;\mathcal{O}_{U'_X})$$

when we write R for $(f')^{-1}i_Y^{-1}\mathcal{O}_{\operatorname{Spec} R}$ where $i_Y : U'_Y \hookrightarrow \operatorname{Spec} R$. Now i_Y is flat so

$$T^i(\mathcal{O}_{U'_X}/R;\mathcal{O}_{U'_X}) \simeq T^i(\mathcal{O}_{U'_X}/(f')^{-1}\mathcal{O}_{U'_Y};\mathcal{O}_{U'_X})$$

and by Lemma 3.2

$$T^i(\mathcal{O}_{U'_X}/(f')^{-1}\mathcal{O}_{U'_Y};\mathcal{O}_{U'_X}) \simeq T^i(\mathcal{O}_{U_X}/f^{-1}\mathcal{O}_{U_Y};\pi_*\mathcal{O}_{U'_X})\,.$$

Therefore after taking invariants and using the conditions we see that

$$T^i(S/R;S)^G \to T^i(\mathcal{O}_{U_X}/f^{-1}\mathcal{O}_{U_Y};\mathcal{O}_{U_X})$$

is an isomorphism for $i = 1$ and injective for $i = 2$. Thus $\mathrm{Def}^G_{S/R} \simeq \mathrm{Hilb}_{U_X/U_Y}$.

We have $\mathcal{H}^i_Z(\mathcal{O}_X) = 0$ for $i \leq 1$ by Lemma 3.3 so again the spectral sequence and vanishing of $T^0(\mathcal{O}_X/f^{-1}\mathcal{O}_Y;\mathcal{O}_X)$ implies $T^i_Z(\mathcal{O}_X/f^{-1}\mathcal{O}_Y;\mathcal{O}_X) = 0$ for $i \leq 2$. We apply the long exact sequence in Theorem 1.1 again to get $T^1(\mathcal{O}_X/f^{-1}\mathcal{O}_Y;\mathcal{O}_X) \simeq T^1(\mathcal{O}_{U_X}/f^{-1}\mathcal{O}_{U_Y};\mathcal{O}_{U_X})$ and $T^2(\mathcal{O}_X/f^{-1}\mathcal{O}_Y;\mathcal{O}_X) \to T^2(\mathcal{O}_{U_X}/f^{-1}\mathcal{O}_{U_Y};\mathcal{O}_{U_X})$ injective. Thus $\mathrm{Hilb}_{X/Y} \simeq \mathrm{Hilb}_{U_X/U_Y}$. □

In the case of the Cox construction for toric varieties (Example 2.5) we get a corollary which is a generalization of the comparison theorem as stated in [PS85]. Proofs and full statements of the results we use here may be found in [CLS11, Chapter 5 and 6].

Let Y be a toric variety with Cox ring R. Every closed subscheme X of Y corresponds to a homogeneous, with respect to the $\mathrm{Cl}(Y)$ grading, ideal $I \subseteq R$. Moreover there is a sheafification construction taking any graded R-module M to a sheaf $\widetilde{M}$ on Y. In particular if $S = R/I$ then $\widetilde{S} = \bigoplus_{\alpha\in \mathrm{Cl}(Y)} \mathcal{O}_X(\alpha)$. Also, as in the case of projective space, one may compute sheaf cohomology from local cohomology at the irrelevant ideal ([EMS00, Proposition 2.3]). In particular there is an exact sequence

$$0 \to H^0_B(S) \to S \to \bigoplus_{\alpha\in \mathrm{Cl}(Y)} H^0(X, \mathcal{O}_X(\alpha)) \to H^1_B(S) \to 0\,.$$

We therefore get the following result for subschemes of toric varieties.

Corollary 3.5. *Let X be a subscheme of a toric variety Y corresponding to the homogeneous ideal I in the Cox ring R of Y. Let Z be the singular locus of Y. Assume I is generated by homogeneous polynomials of degrees $\alpha_1, \dots, \alpha_m \in Cl(Y)$. If*

(i) $\operatorname{depth}_Z \bigoplus_{\alpha \in \mathrm{Cl}(Y)} \mathcal{O}_X(\alpha) \geq 2$

(ii) $(S/I)_{\alpha_i} \simeq H^0(X, \mathcal{O}_X(\alpha_i))$ *for all* $i = 1, \dots m$

then $\mathrm{Def}^0_{S/R}$ *and* $\mathrm{Hilb}_{X/Y}$ *are isomorphic deformation functors.*

Corollary 3.6. *Let $X = \operatorname{Proj} S$ be a subscheme of weighted projective space $\mathbb{P}(\mathbf{q}) = \operatorname{Proj} R$ defined by the homogeneous ideal I. Let $\mathfrak{m}$ be the irrelevant maximal ideal of R and Z the singular locus of $\mathbb{P}(\mathbf{q})$. If*

(i) $\operatorname{depth}_Z \bigoplus_{m \in \mathbb{Z}} \mathcal{O}_X(m) \geq 2$

(ii) $\operatorname{Hom}_R(I, H^1_{\mathfrak{m}}(S))_0 = 0$

then $\mathrm{Def}^0_{S/R}$ *and* $\mathrm{Hilb}_{X/Y}$ *are isomorphic deformation functors.*

3.3 Examples

We give some examples from (straight) projective space to illustrate the condition in Theorem 3.4 and Corollary 3.5. We stick with the above notation so that $Y = \mathbb{P}^n = \operatorname{Proj} R$, $X = \operatorname{Proj} S$, etc. Note here of course Z is empty so only the second condition applies.

Example 3.7. *Points.* If X is s points in general enough position in $\mathbb{P}^n$, then the Hilbert function of X is

$$h_X(\nu) = \inf\{s, \binom{\nu + n}{n}\}$$

by e.g [GMR83]. Let ν_0 be the smallest integer with

$$s \leq \binom{\nu_0 + n}{n}.$$

The exact sequence

$$0 \to S_\nu \to H^0(X, \mathcal{O}_X(\nu)) \to H^1(\mathcal{I}_X(\nu)) \to 0$$

and the fact that $h^0(\mathcal{O}_X(\nu)) = s$ yield $I_\nu = 0$ for $\nu < \nu_0$ and $H^1_{\mathfrak{m}}(S)_\nu = 0$ for $\nu \geq \nu_0$ so $\mathrm{Def}^0_{S/R} \simeq \mathrm{Hilb}_{X/\mathbb{P}^n}$ in this case.

Six general points in $\mathbb{P}^2$ will have Hilbert function $(1, 3, 6, 6, \dots)$. On the other hand the complete intersection of a quadric and a cubic will have $h_X = (1, 3, 5, 6, 6, \dots)$. Thus $\dim I_2 = h^1(\mathcal{I}_X(2)) = 1$, so $\operatorname{Hom}(I, H^1_{\mathfrak{m}}(S))_0 \simeq k$ and the functors need not be isomorphic. Indeed since both obstruction spaces vanish (S is a complete intersection) the map $\mathrm{Def}^0_{S/R} \to \mathrm{Hilb}_{X/\mathbb{P}^n}$ corresponds dually to a surjection of formally smooth complete k-algebras.

Note that we may think of $\mathrm{Def}^0_{S/R}$ as deformations that preserve the Hilbert function and that they correspond to Hilbert function strata of the Hilbert scheme, see [Kle07]. The above two cases show that the Hilbert function strata given by $(1,3,6,6,\dots)$ is an open subscheme of $\mathrm{Hilb}^6(\mathbb{P}^2)$ while $(1,3,5,6,\dots)$ gives a smooth codimension 1 stratum.

Example 3.8. *Curves.* Consider a smooth curve C sitting on a smooth hypersurface V of degree $s \le 3$ in $\mathbb{P}^3$. Let $C = \operatorname{Proj} S \subset \mathbb{P}^3 = \operatorname{Proj} R$ with $S = R/I$. There is an exact sequence of normal bundles

$$0 \to \mathcal{N}_{C/V} \to \mathcal{N}_{C/\mathbb{P}^3} \to \mathcal{O}_C(s) \to 0$$

and $\mathcal{N}_{C/V} \simeq \omega_C \otimes \omega_V^{-1}$. Thus by Serre duality $H^1(\mathcal{N}_{C/V}) = 0$ and $H^1(\mathcal{N}_{C/\mathbb{P}^3}) \simeq H^1(\mathcal{O}_C(s))$ while $H^0(\mathcal{N}_{C/\mathbb{P}^3}) \to H^0(\mathcal{O}_C(s))$ is surjective.

Applying this to the long exact sequence in Theorem 1.1 we get

$$0 \to \mathrm{Hom}_R(I,S)_0 \to H^0(\mathcal{N}_{C/\mathbb{P}^3}) \to \mathrm{Hom}_R(I, H^1_{\mathfrak{m}}(S)) \to T^2(R/S,R)_0 \to 0\,.$$

There is a commutative diagram

$$\begin{tikzcd}
H^0(\mathcal{N}_{C/\mathbb{P}^3}) \arrow[r] \arrow[d, two heads] & \mathrm{Hom}_R(I, H^1_{\mathfrak{m}}(S)) \arrow[d] \\
H^0(\mathcal{O}_C(s)) \arrow[r, two heads] & H^1(\mathcal{I}_C(s))
\end{tikzcd}$$

so if $\mathrm{Hom}_R(I, H^1_{\mathfrak{m}}(S)) \simeq H^1(\mathcal{I}_C(s))$ then $\mathrm{Def}^0_{S/R}$ is unobstructed. If $s = d_0, d_1, \dots, d_m$ are the degrees of the generators of I, then one can show that there is such an isomorphism if $H^1(\mathcal{I}_C(d_i)) = 0$ for all $i = 1, \dots, m$.

If $s = 2$ then $V \simeq \mathbb{P}^1 \times \mathbb{P}^1$ and we can use the Künneth formula to compute these cohomology groups. The outcome is that indeed $\mathrm{Hom}_R(I, H^1_{\mathfrak{m}}(S)) \simeq H^1(\mathcal{I}_C(2))$ so $T^2(R/S,R)_0 = 0$. Moreover if C has bidegree (p,q) with $p \le q$ then

$$h^1(\mathcal{I}_C(2)) = \begin{cases} 0 & \text{if } p > 2 \\ \max\{0, q-3\} & \text{if } p = 2 \end{cases}.$$

Thus $\mathrm{Def}^0_{R/S} \simeq \mathrm{Hilb}_{C/\mathbb{P}^3}$ if $p > 2$ or $q = 3$ while $\mathrm{Def}^0_{R/S}$ corresponds to a smooth stratum of codimension $q-3$ in $\mathrm{Hilb}^{d,g}(\mathbb{P}^3)$ otherwise.

If $s = 3$ then among curves on cubic in $\mathbb{P}^3$ we find the curve that gives rise to Mumford's example of an irreducible component of $\mathrm{Hilb}^{14,24}(\mathbb{P}^3)$ which is not reduced at its generic point ([Mum62]). Let H be a hyperplane section of V and E a line on V. Take C to be a generic element of the complete linear system $|4H + 2E|$ on V. Such a curve may be constructed using linkage in a $(3,6)$ complete intersection and there is a resolution

$$0 \to R(-9) \to R(-8)^2 \oplus R(-7)^2 \to R(-6)^3 \oplus R(-3) \to I \to 0\,.$$

One computes that $h^1(\mathcal{I}_C(d)) = 0$ when $d \ge 6$ and $h^1(\mathcal{I}_C(3)) = h^1(\mathcal{O}_C(3)) = 1$, so $\mathrm{Hom}_R(I, H^1_{\mathfrak{m}}(S)) \simeq H^1(\mathcal{I}_C(3)) \simeq k$ and $T^2(R/S,R)_0 = 0$. Since $\mathrm{Def}^0_{R/S}$ is unobstructed it corresponds to the reduced subscheme of this component.

4 Deformations of the scheme - Def_X

4.1 Results

Lemma 4.1. *If (X, Z, S, J) is a G-ferry quadruple with ferry $\mathcal{S}$ and*

(i) $T^2(\mathcal{S}/\mathcal{O}_X; \mathcal{S})^G = 0$

(ii) $T^2_J(S/k; S)^G = 0$

then $\operatorname{Def}^G_S \to \operatorname{Def}_X$ is smooth. If moreover $T^1(\mathcal{S}/\mathcal{O}_X; \mathcal{S})^G = 0$ and $T^1_J(S/k; S)^G = 0$ then $\operatorname{Def}^G_S \to \operatorname{Def}_X$ is an isomorphism.

Proof. First note that since taking G invariants is exact we have $T^i_X(\mathcal{S})^G \simeq T^i_X$. Secondly [And74, Appendice. Proposition 56] and the fact that π is affine imply $T^i_X(\mathcal{S}) \simeq T^i(\pi^{-1}\mathcal{O}_X/k; \mathcal{O}_{X'})$. The Zariski-Jacobi sequence for $k \to \pi^{-1}\mathcal{O}_X \to \mathcal{O}_{X'}$ reads

$$\to T^i(\mathcal{O}_{X'}/\pi^{-1}\mathcal{O}_X; \mathcal{O}_{X'}) \to T^i_{X'} \to T^i(\pi^{-1}\mathcal{O}_X/k; \mathcal{O}_{X'}) \to T^{i+1}(\mathcal{O}_{X'}/\pi^{-1}\mathcal{O}_X; \mathcal{O}_{X'}) \to$$

so the first condition yields $(T^i_{X'})^G \to T^i_X$ surjective for $i = 1$ and injective for $i = 2$. The exact sequence in Theorem 1.1 for $X' \subseteq \operatorname{Spec} S$ is

$$\cdots \to T^i_J(S/k; S) \to T^i_S \to T^i_{X'} \to T^{i+1}_J(S/k; S) \to \cdots$$

so the second condition implies the same for $(T^i_S)^G \to (T^i_{X'})^G$ and we have proven the first statement. The second statement statement follows from the same argument for $i = 1$. ☐

Even if the second condition above is not satisfied the construction can help computing deformations.

Lemma 4.2. *If (X, Z, S, J) is a G-ferry quadruple with ferry $\mathcal{S}$ and*

(i) $T^1(\mathcal{S}/\mathcal{O}_X; \mathcal{S})^G = T^2(\mathcal{S}/\mathcal{O}_X; \mathcal{S})^G = 0$

(ii) $(T^1_S)^G = (T^2_S)^G = 0$

then $T^1_X \simeq T^2_J(S/k; S)^G$ and an injective map $T^2_X \hookrightarrow T^3_J(S/k; S)^G$.

Proof. This follows from the exact sequences in the proof of Lemma 4.1. ☐

Lemma 4.3. *If (X, Z, S, J) is a G-ferry quadruple then $T^1_J(S/k; S) \simeq \operatorname{Der}_k(S, H^1_J(S))$ and there is an exact sequence*

$$0 \to T^1(S/k; H^1_J(S)) \to T^2_J(S/k; S) \to T^0(S/k; H^2_J(S)) \to T^2(S/k; H^1_J(S))\,.$$

In particular if $\operatorname{depth}_J S \geq 2$ then $T^2_J(S/k; S) \simeq \operatorname{Der}_k(S, H^2_J(S))$ as SG-modules.

Proof. Since $H^0_J(S) = 0$ the exact sequence comes from the edge exact sequence for the spectral sequence in Theorem 1.1. ☐

Lemma 4.4. *If (X, Z, S, J) is a G-ferry quadruple and* $\operatorname{depth}_Z \mathcal{S} \geq 2$ *then*

$$\mathcal{T}^0(\mathcal{S}/\mathcal{O}_X; \mathcal{S}) \simeq \mathcal{S} \otimes_k \mathfrak{g}$$

where $\mathfrak{g}$ is the Lie algebra of G.

Proof. Recall that $\pi_{|U'}$ is a locally trivial G-bundle, and $\Theta_G = \mathcal{O}_G \otimes_k \mathfrak{g}$. The group action gives a global trivialization so $\Theta_{U'/U} \simeq \mathcal{O}_{U'} \otimes_k \mathfrak{g}$. Since $\mathcal{T}^0(\mathcal{S}/\mathcal{O}_X; \mathcal{S})$ is the dual of $\Omega^1_{\mathcal{S}/\mathcal{O}_X}$ it also has depth ≥ 2 in Z. It is therefore isomorphic to $i_*\pi_*\Theta_{U'/U} \simeq \mathcal{S} \otimes_k \mathfrak{g}$. ☐

Definition 4.5. Assuming $\operatorname{depth}_Z \mathcal{S} \geq 2$, let $E_1, \dots, E_r$ be a set of invariant global sections which generate the free sheaf $\mathcal{T}^0(\mathcal{S}/\mathcal{O}_X; \mathcal{S})$. We call such a set a *set of Euler derivations* for the G-ferry quadruple. A set of Euler derivations defines an $\mathcal{S}G$ map $E : \Omega^1_{\mathcal{S}/\mathcal{O}_X} \to \mathcal{S}^r$. Let $\mathcal{Q} = \operatorname{coker}(E)$. Let $Z(\mathcal{Q})$ be the support of $\mathcal{Q}$.

Remark. The morphism E is just the natural map $\Omega^1_{\mathcal{S}/\mathcal{O}_X} \to (\Omega^1_{\mathcal{S}/\mathcal{O}_X})^{\vee\vee}$ after choosing a basis for the free sheaf $(\Omega^1_{\mathcal{S}/\mathcal{O}_X})^{\vee\vee}$ thus $\ker(E)$ is the torsion submodule of $\Omega^1_{\mathcal{S}/\mathcal{O}_X}$ and $\mathcal{Q}$ is independent of the choice of Euler derivations. Note that $Z(\mathcal{Q}) \subseteq Z$.

Remark. If G is finite and $\operatorname{depth}_Z \mathcal{S} \geq 2$ then $\mathcal{T}^0(\mathcal{S}/\mathcal{O}_X; \mathcal{S})$ vanishes so $\mathcal{T}or(\Omega^1_{\mathcal{S}/\mathcal{O}_X}) = \Omega^1_{\mathcal{S}/\mathcal{O}_X}$. Thus $Z(\mathcal{Q})$ contains the closed locus where *the isotropy group of the G action on X' is finite.* For toric varieties in fact $Z(\mathcal{Q})$ equals the non-simplicial locus of X.

Lemma 4.6. *If (X, Z, S, J) is a G-ferry quadruple with* $\operatorname{depth}_Z \mathcal{S} \geq 2$ *then*

$$\mathcal{T}^i(\mathcal{S}/\mathcal{O}_X; \mathcal{S}) \simeq \mathcal{E}xt^i_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S}) \text{ for } i = 0, 1, 2.$$

Proof. Consider the spectral sequence $\mathcal{E}xt^p_{\mathcal{S}}(\mathcal{T}_q(\mathcal{S}/\mathcal{O}_X; \mathcal{S}), \mathcal{S}) \Rightarrow \mathcal{T}^{p+q}(\mathcal{S}/\mathcal{O}_X; \mathcal{S})$. Since the $\mathcal{T}_q(\mathcal{S}/\mathcal{O}_X; \mathcal{S})$ are supported on Z when $q \geq 1$, the depth condition yields the result. ☐

Assume $\mathcal{S}$ is a sheaf of $\mathcal{O}_X$-algebras on a scheme X and $Z \subseteq X$ is a locally closed. If $\mathcal{F}$ and $\mathcal{G}$ are $\mathcal{S}$-modules, denote by $\operatorname{Ext}^i_{\mathcal{S},Z}(\mathcal{F}, \mathcal{G})$, respectively $\mathcal{E}xt^i_{\mathcal{S},Z}(\mathcal{F}, \mathcal{G})$, the higher derived functors of $\mathcal{G} \mapsto H^0_Z(X, \mathcal{H}om_{\mathcal{S}}(\mathcal{F}, \mathcal{G}))$, respectively $\mathcal{G} \mapsto \mathcal{H}^0_Z(X, \mathcal{H}om_{\mathcal{S}}(\mathcal{F}, \mathcal{G}))$. We refer to SGA 2 Exposè VI ([Gro05]) for details and the results we will use.

Lemma 4.7. *If (X, Z, S, J) is a G-ferry quadruple with* $\operatorname{depth}_Z \mathcal{S} \geq 2$ *and* $\operatorname{depth}_Z \mathcal{O}_X \geq 3$ *then $\mathcal{E}xt^1_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})^G = 0$ and*

$$T^i(\mathcal{S}/\mathcal{O}_X; \mathcal{S})^G \simeq \operatorname{Ext}^i_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})^G \text{ for } i = 0, 1, 2\,.$$

Proof. By Lemma 4.4, $\Omega^1_{\mathcal{S}/\mathcal{O}_X|U} \simeq \mathcal{S}^r_{|U}$ so we get an exact sequence

$$\to \mathcal{E}xt^1_{\mathcal{S},Z}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S}) \to \mathcal{E}xt^1_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S}) \to \mathcal{R}^1 j_* \mathcal{S}^r_{|U} \to$$

from the long exact sequence for $\mathcal{E}xt^i_{\mathcal{S},Z}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})$. Now since $\operatorname{depth}_Z \mathcal{S} \geq 2$ the left sheaf vanishes. The right sheaf is isomorphic to $r\mathcal{H}^2_Z(\mathcal{S})$. After taking invariants this becomes $r\mathcal{H}^2_Z(\mathcal{O}_X)$ which also vanishes since $\operatorname{depth}_Z \mathcal{O}_X \geq 3$. Thus $\mathcal{E}xt^1_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})^G = 0$.

We have local global spectral sequences which at the E_2 level are

$$\begin{gathered} H^p(X, \mathcal{T}^q(\mathcal{S}/\mathcal{O}_X;\mathcal{S})) \Rightarrow T^{p+q}(\mathcal{S}/\mathcal{O}_X;\mathcal{S}) \\ H^p(X, \mathcal{E}xt^q_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})) \Rightarrow \mathrm{Ext}^{p+q}_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S}) \end{gathered}$$

Lemma 4.6 implies the isomorphisms for $i = 0, 1$. But since $\mathcal{E}xt^1_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})^G = 0$ the spectral sequences yield exact sequences

$$\begin{gathered} 0 \to H^2(X, \mathcal{T}^0(\mathcal{S}/\mathcal{O}_X;\mathcal{S}))^G \to T^2(\mathcal{S}/\mathcal{O}_X;\mathcal{S})^G \to H^0(X, \mathcal{T}^2(\mathcal{S}/\mathcal{O}_X;\mathcal{S}))^G \to 0 \\ 0 \to H^2(X, \mathcal{E}xt^0_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S}))^G \to \mathrm{Ext}^2_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})^G \to H^0(X, \mathcal{E}xt^2_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S}))^G \to 0 \end{gathered}$$

so the isomorphism for $i = 2$ also follows from Lemma 4.6. □

Proposition 4.8. *If (X, Z, S, J) is a G-ferry quadruple with*

(i) $\mathrm{depth}_Z\, \mathcal{S} \geq 2$

(ii) $\mathrm{depth}_Z\, \mathcal{O}_X \geq 3$

then $T^1(\mathcal{S}/\mathcal{O}_X;\mathcal{S})^G \simeq rH^1(X, \mathcal{O}_X)$ and there are exact sequences

$$\begin{gathered} 0 \to rH^2(X, \mathcal{O}_X) \to T^2(\mathcal{S}/\mathcal{O}_X;\mathcal{S})^G \to H^0(X, \mathcal{E}xt^2_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})^G) \to 0 \\ 0 \to \mathrm{Hom}_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{H}^2_Z(\mathcal{S}))^G \to H^0(X, \mathcal{E}xt^2_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})^G) \to \mathrm{Hom}_{\mathcal{S}}(\mathcal{Q}, \mathcal{H}^3_Z(\mathcal{S}))^G . \end{gathered}$$

Moreover there are inclusions $\mathrm{Hom}_{\mathcal{S}}(\mathcal{Q}, \mathcal{H}^3_Z(\mathcal{S}))^G \subseteq \mathrm{Hom}_{\mathcal{O}_X}(\mathcal{Q}^G, \mathcal{H}^3_Z(\mathcal{O}_X)) \subseteq rH^3_Z(\mathcal{O}_X)$. *If $\mathcal{S}$ is regular then the second sequence is exact on the right as well.*

Proof. The first statement and the first exact sequence follow from the local-global spectral sequence and Lemma 4.4 and 4.7. Let j be the inclusion of U in X. From the long exact sequence for $\mathcal{E}xt^i_{\mathcal{S},Z}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})$ we get

$$\to \mathcal{R}^1 j_* \mathcal{S}^r_{|U} \to \mathcal{E}xt^2_{\mathcal{S},Z}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S}) \to \mathcal{E}xt^2_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S}) \to \mathcal{R}^2 j_* \mathcal{S}^r_{|U} \to \mathcal{E}xt^3_{\mathcal{S},Z}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S}) .$$

Taking invariants and using the depth assumption this becomes

$$0 \to \mathcal{E}xt^2_{\mathcal{S},Z}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})^G \to \mathcal{E}xt^2_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})^G \to r\mathcal{H}^3_Z(\mathcal{O}_X) \to \mathcal{E}xt^3_{\mathcal{S},Z}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})^G .$$

Using the spectral sequence $\mathcal{E}xt^p_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{H}^q_Z(\mathcal{S})) \Rightarrow \mathcal{E}xt^{p+q}_{\mathcal{S},Z}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})$ and the conditions this becomes again

$$0 \to \mathcal{E}xt^0_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{H}^2_Z(\mathcal{S}))^G \to \mathcal{E}xt^2_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})^G \to r\mathcal{H}^3_Z(\mathcal{O}_X) \xrightarrow{u} \mathcal{E}xt^3_{\mathcal{S},Z}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S})^G .$$

There is a composite map $\varepsilon = v \circ u$

$$r\mathcal{H}^3_Z(\mathcal{S}) \xrightarrow{u} \mathcal{E}xt^3_{\mathcal{S},Z}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{S}) \xrightarrow{v} \mathcal{H}om_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{H}^3_Z(\mathcal{S}))$$

where v is the map to $E_2^{0,3}$ in the above spectral sequence. Thus $\ker u \subseteq \ker \varepsilon$. On the other hand ε is the map induced by the middle map in the exact sequence

$$0 \to \mathcal{T}or(\Omega^1_{\mathcal{S}/\mathcal{O}_X}) \to \Omega^1_{\mathcal{S}/\mathcal{O}_X} \to \mathcal{S}^r \to \mathcal{Q} \to 0 \tag{4.1}$$

from Definition 4.5. Therefore $\ker \varepsilon = \mathcal{H}om_{\mathcal{S}}(\mathcal{Q}, \mathcal{H}^3_Z(\mathcal{S}))$ and we have shown that the second sequence in the statement is exact.

The sequence (4.1) is an $\mathcal{S}G$ exact sequence so $\mathcal{Q}$ is generated by G invariants. Thus $\mathcal{H}om_{\mathcal{S}}(\mathcal{Q}, \mathcal{H}^3_Z(\mathcal{S}))^G \to \mathcal{H}om_{\mathcal{O}_X}(\mathcal{Q}^G, \mathcal{H}^3_Z(\mathcal{S})^G)$ is injective. The second inclusion follows from first taking invariants of (4.1) and then applying $\mathcal{H}om(-, \mathcal{H}^3_Z(\mathcal{O}_X))$.

To prove the last statement note that if $\mathcal{S}$ is regular then locally $\mathcal{O}_X \to \mathcal{S}$ is a localization of some $A^G \to A$ as in Lemma 1.3. Therefore $\mathcal{E}xt^1_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{H}^2_Z(\mathcal{S}))^G = 0$ and the map v above is injective as well. It follows that $\ker u = \ker \varepsilon$. ☐

Example 4.9. *Finite G.* We can apply the above to the situation in Example 2.7, i.e. $G \subset \mathrm{GL}_n(\mathbb{C})$ is finite without pseudo-reflections and

$$(X, Z, S, J) = (\mathbb{C}^n/G, \mathrm{Sing}(\mathbb{C}^n/G), \mathcal{O}_{\mathbb{C}^n}, (1))\,.$$

Assume that $\mathrm{codim}_Z X \geq 3$ (the singularity need not be isolated). Then since X is Cohen-Macaulay $\mathrm{depth}_Z \mathcal{O}_X \geq 3$ and since G is finite $\mathrm{depth}_Z S \geq 3$. We also have $r = 0$ so it follows from Proposition 4.8, Lemma 4.6 and Lemma 4.1 that X is rigid. This generalizes Schlessinger's result in [Sch71].

In general consider an affine G-ferry quadruple

$$(X, Z, S, J) = (\mathrm{Spec}\, S^G, Z, S, (1))$$

with finite G and Z the locus where $\pi : \mathrm{Spec}\, S \to X$ is not étale. Then by the above results we have $\mathrm{Def}^G_S \simeq \mathrm{Def}_X$ if $\mathrm{depth}_Z S \geq 3$. If particular if S is Cohen-Macaulay then it is enough to assume $\mathrm{codim}\, Z \geq 3$. See also [Ste88, Section 7].

Theorem 4.10. *If (X, Z, S, J) is a G-ferry quadruple with*

(i) $\mathrm{depth}_Z \mathcal{S} \geq 2$ *and* $\mathrm{depth}_Z \mathcal{O}_X \geq 3$

(ii) *G is finite or* $H^2(X, \mathcal{O}_X) = 0$

(iii) $H^0(X, \mathcal{E}xt^2_{\mathcal{S}}(\Omega^1_{S/\mathcal{O}_X}, \mathcal{S})^G) = 0$

(iv) $T^1(S/k; H^1_J(S))^G = 0$ *and* $\mathrm{Der}_k(S, H^2_J(S))^G = 0$

then $\mathrm{Def}^G_S \to \mathrm{Def}_X$ *is smooth. If moreover*

(i) *G is finite or* $H^1(X, \mathcal{O}_X) = 0$

(ii) $\mathrm{Der}_k(S, H^1_J(S))^G = 0$

then $\mathrm{Def}^G_S \to \mathrm{Def}_X$ *is an isomorphism.*

Proof. The result follows directly from Lemma 4.1 and Proposition 4.8. ☐

In low dimensional cases, we can not expect that $\mathrm{Def}^G_S \to \mathrm{Def}_X$ is an isomorphism, even though the first four assumptions of the theorem are satisfied. Here is an example of space curves.

Example 4.11. For space curves we only need to verify (iv) to conclude that $\mathrm{Def}^0_S \to \mathrm{Def}_X$ is smooth. We claim that a general space curve $X = \operatorname{Proj} S$ in $\mathrm{Hilb}^{d,g}(\mathbb{P}^3)$ satisfying $g < d+3$ also satisfy (iv). Indeed in this case J is the irrelevant maximal ideal and $H^2_J(S) \simeq \sum_\nu H^1(X, O_X(\nu))$. We get $\mathrm{Der}_k(S, H^2_J(S))_0 = 0$ provided $H^1(X, \mathcal{O}_X(1)) = 0$. It is well known that a general curve satisfies this property.

We need to show $T^1(S/k, H^1_J(S))_0 = 0$. Let I be the homogeneous ideal of X in $\mathbb{P}^3 = \operatorname{Proj} R$. Since $\mathrm{Hom}_R(I, H^1_J(S))_0 = T^1(S/R, H^1_J(S)) \to T^1(S/k, H^1_J(S))_0$ is surjective, it suffices to show $\mathrm{Hom}(I, H^1_J(S))_0 = 0$. Space curves of maximal rank (i.e. $H^1(\mathcal{I}_X(v)) = 0$ provided $H^0(\mathcal{I}_X(v)) \neq 0$) necessarily satisfy this property. The main theorem Ellia and Ballico in [BE85] implies that the general curve in the range $g < d-3$ has maximal rank. It follows that $\mathrm{Def}^0_S \to \mathrm{Def}_X$ is smooth in this range. But if $g > 0$ then $H^1(X, \mathcal{O}_X) \neq 0$ and we do not expect $\mathrm{Def}^0_S \to \mathrm{Def}_X$ to be an isomorphism.

4.2 The Cohen-Macaulay case

Since we are dealing so much with depth conditions we get better results when X is Cohen-Macaulay.

Proposition 4.12. *If (X, Z, S, J) is a G-ferry quadruple with*

(i) *X an integral Cohen-Macaulay scheme*

(ii) $\mathrm{depth}_Z \mathcal{S} \geq 2$

(iii) $\operatorname{codim} Z \geq 3$

(iv) $\operatorname{codim} Z(\mathcal{Q}) \geq 4$

then there is an exact sequence

$$0 \to rH^2(X, \mathcal{O}_X) \to T^2(\mathcal{S}/\mathcal{O}_X; \mathcal{S})^G \to \mathrm{Hom}_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{H}^2_Z(\mathcal{S}))^G \to 0$$

Proof. We will prove that $\mathcal{H}om_{\mathcal{O}_X}(\mathcal{Q}^G, \mathcal{H}^3_Z(\mathcal{O}_X)) = 0$ and then the result follows from Proposition 4.8. But

$$\mathcal{H}om_{\mathcal{O}_X}(\mathcal{Q}^G, \mathcal{H}^3_Z(\mathcal{O}_X)) = \mathcal{H}om_{\mathcal{O}_X}(\mathcal{Q}^G, \mathcal{H}^0_{Z(\mathcal{Q})}(\mathcal{H}^3_Z(\mathcal{O}_X))) = 0$$

because of the assumptions and Lemma 1.5. ☐

Theorem 4.13. *If (X, Z, S, J) is a G-ferry quadruple with X an integral Cohen-Macaulay scheme and*

(i) $\operatorname{depth}_Z \mathcal{S} \geq 2$, $\operatorname{codim} Z \geq 3$ *and* $\operatorname{codim} Z(\mathcal{Q}) \geq 4$

(ii) *G is finite or* $H^2(X, \mathcal{O}_X) = 0$

(iii) $\operatorname{Hom}_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{H}^2_Z(\mathcal{S}))^G = 0$

(iv) $T^1(S/k; H^1_J(S))^G = 0$ *and* $\operatorname{Der}_k(S, H^2_J(S))^G = 0$

then $\operatorname{Def}^G_S \to \operatorname{Def}_X$ *is smooth. If moreover*

(i) *G is finite or* $H^1(X, \mathcal{O}_X) = 0$

(ii) $\operatorname{Der}_k(S, H^1_J(S))^G = 0$

then $\operatorname{Def}^G_S \to \operatorname{Def}_X$ *is an isomorphism.*

Proof. Apply Proposition 4.12. □

Remark. Let J' be the radical ideal of the closure of Z' in $\operatorname{Spec} S$. If $\operatorname{Spec} S$ is smooth and $\operatorname{codim} J' \geq 2$ then $\operatorname{Der}_k(S, H^2_J(S))^G \simeq T^2_J(S/k; S)^G \simeq (T^1_{X'})^G \simeq H^1(X', \Theta_{X'})^G \simeq H^1(X, \Theta_X)$. This means that $\operatorname{Der}_k(S, H^2_J(S))^G$ in this case computes the locally trivial first order deformations.

By the Hochster-Roberts theorem X will be Cohen-Macaulay if S is a regular ring. In that case we get a statement about rigidity of X.

Corollary 4.14. *Assume (X, Z, S, J) is a G-ferry quadruple with S a regular ring. Let J' be the radical ideal of the closure of Z' in* $\operatorname{Spec} S$. *If*

(i) $\operatorname{codim} J' \geq 2$, $\operatorname{codim} Z \geq 3$ *and* $\operatorname{codim} Z(\mathcal{Q}) \geq 4$

(ii) *G is finite or* $H^2(X, \mathcal{O}_X) = 0$

(iii) $\operatorname{Hom}_{\mathcal{S}}(\Omega^1_{\mathcal{S}/\mathcal{O}_X}, \mathcal{H}^2_Z(\mathcal{S}))^G = 0$

(iv) $\operatorname{Der}_k(S, H^2_J(S))^G = 0$

then X is rigid.

4.3 Applications to toric varieties

Applied to toric varieties the above would only give rigidity results. One key object to understand is $H^2_B(S)$. Let $\{m_1, \dots, m_r\}$ be monomial generators for any squarefree monomial ideal $B \subseteq S$. For $I \subseteq \{1, \dots, r\}$ let T_I be the simplicial complex on the vertex set $\{1, \dots, r\}$ where $\{j_1, \dots, j_k\}$ is a face if $x_i \nmid \operatorname{lcm}(m_{j_1}, \dots, m_{j_k})$ for some $i \in I$. If $p \in \mathbb{Z}^N$ define $\operatorname{neg}(p) \subseteq \{1, \dots, N\}$ to be the set $\{i \mid p_i \leq -1\}$.

Theorem 4.15. ([Mus00, Theorem 2.1]) *If $p \in \mathbb{Z}^N$ then there are isomorphisms $H^i_B(S)_p \simeq \widetilde{H}^{i-2}(T_I; k)$ when $I = \operatorname{neg}(p)$. Moreover the map*

$$H^i_B(S)_p \xrightarrow{\cdot x_i} H^i_B(S)_{p+e_i}$$

corresponds to the map $H^{i-2}(T_{\operatorname{neg}(p)}; k) \to H^{i-2}(T_{\operatorname{neg}(p+e_i)}; k)$ induced in cohomology by the inclusion $T_{\operatorname{neg}(p+e_i)} \subseteq T_{\operatorname{neg}(p)}$. In particular, if $p_i \neq -1$, then $\cdot x_i$ is an isomorphism.

From now on assume S is the Cox ring and B the irrelevant ideal for a fan Σ.

Lemma 4.16. *The codimension 2 prime ideals of B are the (x_i, x_j) with ρ_i and ρ_j not in the same cone in Σ.*

Proof. This follows directly from the description of the prime components of B. See e.g. [CLS11, Proposition 5.1.6]. ☐

Let B_2 be the intersection of the codimension 2 primes of B. Let K be the simplicial complex which has B_2 as Stanley-Reisner ideal. Let Γ be the graph with vertices $\{0, \dots, N\}$ and edges $\{i, j\}$ when ρ_i and ρ_j are in the same cone in Σ. Define $C(\Gamma)$ to be the clique complex of Γ. Let Γ_I be the induced subgraph with vertices in I.

Lemma 4.17. *The complex K is the Alexander dual of $C(\Gamma)$. In particular there is a one to one correspondance between facets of $C(\Gamma)$ and a minimal generating set for B_2 given by $F \mapsto x_{F^c}$. This correspondance identifies T_I with the complex with vertex set equal the set of facets of $C(\Gamma)$ containing an element of I and $\{F_{i_1}, \dots, F_{i_k}\}$ is a face if $i \in \bigcap F_{i_j}$ for some $i \in I$.*

Proof. The facets of K are the complements $\{i, j\}^c$ where $\{i, j\}$ is a non-edge of Γ. We have $f \in C(\Gamma)^\vee \Leftrightarrow f^c \notin C(\Gamma) \Leftrightarrow f^c$ contains a non-edge of $\Gamma \Leftrightarrow f \subseteq \{i, j\}^c$ for a non-edge $\{i, j\} \Leftrightarrow f \in K$. A set $\{x_{F^c_{i_1}}, \dots, x_{F^c_{i_k}}\}$ is a face of T_I if there is an $i \in I$ with $i \notin \bigcup F^c_{i_k} = (\bigcap F_{i_k})^c$, that is if $i \in \bigcap F_{i_k}$. ☐

Lemma 4.18. *If $i \in I$ let f_i be the face of T_I given by $\{F : i \in F\}$. The map $H_0(\Gamma_I) \to H_0(T_I)$ defined by $[i] \mapsto [F]$ where F is any vertex in f_i is an isomorphism.*

Proof. First note that $\{i, j\}$ is an edge in Γ_I iff there is a maximal clique F of Γ containing $\{i, j\}$ which is iff $f_i \cap f_j \neq \emptyset$. To show that the map is well defined assume $[i] = [j]$. Then there is an edge path through vertices $i = i_0, i_1, \dots, i_r = j$ in Γ_I. Then since the $f_{i_k} \cap f_{i_{k+1}} \neq \emptyset$ we may find a path from any vertex of f_i to any vertex in f_j.

The map is clearly surjective. To show injectivity assume $F \in f_i$ and $G \in f_j$ are connected by an edge path $F = F_0, F_1, \dots, F_r = G$. This means that the intersection of cliques $F_k \cap F_{k+1}$ contains an element $i_k \in I$. This implies that there are edges $\{i_k, i_{k+1}\} \in \Gamma$ and therefore in the induced subgraph Γ_I. Thus $[i] = [j]$. ☐

Proposition 4.19. *There is an isomorphism $H^2_B(S)_p \simeq \widetilde{H}^0(\Gamma_I, k)$ where $I = \operatorname{neg}(p)$.*

Proof. This follows now directly from Theorem 4.15, Lemma 4.17 and 4.18. $\square$

We wish to apply the above with Corollary 4.14 to get rigidity results for toric singularities. If σ is a cone defining an affine toric variety X with Cox ring $S = k[x_1, \dots, x_N]$, let Σ be the fan consisting of smooth faces of σ and $B = B(\Sigma)$ the corresponding irrelevant ideal. If $X = \operatorname{Spec} A$ we must at least have $\operatorname{Hom}_S(\Omega^1_{S/A}, H^2_B(S))_0 = \operatorname{Der}_A(S, H^2_B(S))_0 = 0$ in the $\operatorname{Cl}(X)$ grading. This would follow if $H^2_B(S)_{\alpha_i} = 0$ for $i = 1, \dots, N$ where α_i is the degree of x_i.

Lifting this to the $\mathbb{Z}^N$ grading we need $H^2_B(S)_p = 0$ (with the notation of Example 2.5) when $p = q + e_i$ and $q = (\dots, \langle u, v_j \rangle, \dots)$ for some $u \in M$. If we assume $\operatorname{codim} \operatorname{Sing}(X) \geq 3$ then $H^2_B(S)_0 = H^2_{\operatorname{Sing}(X)}(A) = 0$. In the $\mathbb{Z}^N$ grading this means $H^2_B(S)_q = 0$ when q is as above. Thus by Theorem 4.15 it is enough to check when in q the coordinate $\langle u, v_i \rangle = -1$.

Let Γ^f be the graph with vertices $\{0, \dots, N\}$ and edges $\{i, j\}$ when $\{\rho_i, \rho_j\}$ span a 2-dimensional face of σ and Γ^f_I the induced subgraph as above. If $\operatorname{codim} \operatorname{Sing}(X) \geq 3$ then Γ^f_I is a subgraph of $\Gamma_I(\Sigma)$ with the same vertex set so the map $\widetilde{H}_0(\Gamma^f_I) \to \widetilde{H}_0(\Gamma_I)$ is surjective.

If $u \in M$ and $\langle u, v_i \rangle = -1$ let

$$I = \{j \in \{1, \dots, N\} : \langle u, v_j \rangle \leq -1\} \setminus \{i\}$$

and set $\Gamma_i(u) = \Gamma^f_I$. (These graphs also appear in the study of deformations of smooth complete toric varieties in [Ilt11].) The above arguments together with Corollary 4.14 yield

Proposition 4.20. *If X is an affine toric variety and*

(i) *X is smooth in codimension 2 and simplicial in codimension 3*

(ii) *$\Gamma_i(u)$ is connected for all $u \in M$*

then X is rigid.

Lemma 4.21. *Let P be a polytope, H^* a closed half-space with bounding hyperplane H and v a vertex of P with $v \in H$. If $\Gamma_v(H)$ is the induced subgraph of the edge graph of P on the vertices in*

$$\{w \in \operatorname{vert} P : w \in H^*\} \setminus \{v\}$$

then $\Gamma_v(H)$ is connected.

Proof. If $H \cap \operatorname{int} P \neq \emptyset$ let H' be the supporting hyperplane in H^* which is parallell to H. By standard arguments every vertex in $\Gamma_v(H)$ is either on H' or has a neighbor vertex in H^* which is nearer to H'. Thus $\Gamma_v(H)$ is connected in this case.

If $H \cap \operatorname{int} P = \emptyset$ then H is a supporting hyperplane. If $P \subset H^*$ then $\Gamma_v(H)$ is the edge graph of P with one vertex removed. Since edge graphs of d-polytopes are d-connected $\Gamma_v(H)$ is connected (trivially also for $d = 0, 1$). If $P \cap H^* = F$ is a face then the same argument applied to F yields the result. $\square$

With this we can reprove a theorem of Altmann in [Alt95].

Corollary 4.22. *A $\mathbb{Q}$-Gorenstein affine toric variety which is smooth in codimension 2 and simplicial in codimension 3 is rigid.*

Proof. An affine toric variety is $\mathbb{Q}$-Gorenstein if and only if there is a primitive $u_0 \in M$ and a positive integer g such that $\langle u_0, v_i \rangle = g$ for all generators v_i of the rays of σ. This means that if P is the convex hull of the v_i in the hyperplane $\langle u_0, - \rangle = g$, then for all $u \in M$ the graph $\Gamma_i(u)$ is a $\Gamma_v(H)$ as in Lemma 4.21. ☐

With the same arguments we can reprove Totaro's generalization, [Tot12, Theorem 5.1] of theorems of Bien-Brion and de Fernex-Hacon.

Corollary 4.23. *A toric Fano variety which is smooth in codimension 2 and simplicial in codimension 3 is rigid.*

Proof. Corollary 4.22 takes care of the local situation so what is left to prove is condition (iv) in Corollary 4.14. The Fano condition implies that the fan Σ of a Fano toric variety is the face fan of a polytope. Thus the same arguments as above show that $H^2_{B(\Sigma)}(S)_{\alpha_i} = 0$ where α_i are degrees of the generators of the Cox ring of Σ. ☐

References

[Alt95] Klaus Altmann, *Minkowski sums and homogeneous deformations of toric varieties*, Tohoku Math. J. (2) **47** (1995), no. 2, 151–184.

[And74] Michel André, *Homologie des algèbres commutatives*, Springer-Verlag, 1974.

[BE85] E. Ballico and Ph. Ellia, *The maximal rank conjecture for nonspecial curves in* $\mathbf{P}^3$, Invent. Math. **79** (1985), no. 3, 541–555.

[CLS11] David A. Cox, John B. Little, and Henry K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, 2011.

[Cox95] David A. Cox, *The homogeneous coordinate ring of a toric variety*, J. Algebraic Geom. **4** (1995), no. 1, 17–50.

[Del75] Charles Delorme, *Espaces projectifs anisotropes*, Bull. Soc. Math. France **103** (1975), no. 2, 203–223.

[Dol82] Igor Dolgachev, *Weighted projective varieties*, Group actions and vector fields (Vancouver, B.C., 1981), Lecture Notes in Math., vol. 956, Springer, Berlin, 1982, pp. 34–71.

[EMS00] David Eisenbud, Mircea Mustaţă, and Mike Stillman, *Cohomology on toric varieties and local cohomology with monomial supports*, J. Symbolic Comput. **29** (2000), no. 4-5, 583–600.

[GMR83] A. V. Geramita, P. Maroscia, and L. G. Roberts, *The Hilbert function of a reduced k-algebra*, J. London Math. Soc. (2) **28** (1983), no. 3, 443–452.

[Gro05] Alexander Grothendieck, *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*, Documents Mathématiques (Paris) [Mathematical Documents (Paris)], 4, Société Mathématique de France, Paris, 2005.

[Ilt11] Nathan Owen Ilten, *Deformations of smooth toric surfaces*, Manuscripta Math. **134** (2011), no. 1-2, 123–137.

[Kle79] Jan O. Kleppe, *Deformations of graded algebras*, Math. Scand. **45** (1979), 205–231.

[Kle07] Jan O. Kleppe, *Families of Artinian and one-dimensional algebras*, J. Algebra **311** (2007), no. 2, 665–701.

[Lau79] Olav Arnfinn Laudal, *Formal moduli of algebraic structures*, Lecture Notes in Mathematics, vol. 754, Springer-Verlag, 1979.

[Mum62] David Mumford, *Further pathologies in algebraic geometry*, Amer. J. Math. **84** (1962), 642–648.

[Mus00] Mircea Mustaţă, *Local cohomology at monomial ideals*, J. Symbolic Comput. **29** (2000), no. 4-5, 709–720.

[PS85] Ragni Piene and Michael Schlessinger, *On the Hilbert scheme compactification of the space of twisted cubics*, Amer. J. Math. **107** (1985), 761–774.

[Rim80] Dock S. Rim, *Equivariant G-structure on versal deformations*, Trans. Amer. Math. Soc. **257** (1980), no. 1, 217–226.

[Sch71] Michael Schlessinger, *Rigidity of quotient singularities*, Invent. Math. **14** (1971), 17–26.

[Ste88] J. Stevens, *On canonical singularities as total spaces of deformations*, Abh. Math. Sem. Univ. Hamburg **58** (1988), 275–283.

[Tot12] Burt Totaro, *Jumping of the nef cone for Fano varieties*, J. Algebraic Geom. **21** (2012), no. 2, 375–396.

JAN ARTHUR CHRISTOPHERSEN
MATEMATISK INSTITUTT
POSTBOKS 1053 BLINDERN
UNIVERSITY OF OSLO
N-0316 OSLO, NORWAY

E-mail address: christop@math.uio.no

Jan O. Kleppe
Oslo and Akershus University College
of Applied Sciences
Faculty of Technology, Art and Design
PB 4 St. Olavs plass
N-0130 Oslo, NOrway

E-mail address: JanOddvar.Kleppe@hioa.no